## 5.0 BIBLIOGRAPHY


1. Chein YW. Novel drug delivery systems. 3rd ed, New York, Marcel Dekker, 2002; 139-96.
2. Rawlins EA. Bentley's text book of pharmaceutics. 8th ed. London, Bailliere Tindall, 1992.
3. Leon L, Herbert AL, Josheph LK. The theory and practice of industrial pharmacy. 3rd ed. Bombay, Varghese Publishing House, 1991.
4. Verma RK, Garg S (2001), Current status of drug delivery technologies and future directions. Pharma Techno, 2001: 25(2): 1-13.
5. Leon L, Herbert AL. Compression coated and layer tablets. In: Pharmaceutical Dosage Forms. New York, Marcel Dekker Inc, 2002; 1(2): 247-84.
6. Gennaro AR. Remington, the Science and Practice of Pharmacy. New Delhi, Elsevier, 2000; 20: 903-07.
7. Ravi KN, Sachin RP, Mrithyunjaya BP, Mahalaxmi B. Evaluation of Disintegrating Properties of *Mangifera indica* gum. RJPS, 2011; 1 (1): 11-22.
8. Ravi KN, Narayana SVB, Mahalaxmi R. exploitation of *mangifera indica* gum as novel natural gelling agent in the designing of gel formulations. Pharmacologyonline, 2011; 2: 522-541.
9. Raju M, Sundaramoorthy S, Vetrichelvan T. Formulation and *in-vitro* evaluation of sustained release matrix tablets of aceclofenac by using different natural polymers. RJPBCS, 2010; 1(4): 279-91.
10. Shingala VK, Singh AK, Yadav SK, Sivakumar T. Design and characterization of Diclofenac sodium tablets containing *Mangifera indica* resin as release retardant. Int J Pharm Tech Res, 2010; 2(3): 2108-11.
11. Velmurugan S, Deepika B, Nagaraju K, Vinushitha S. Formulation and *in-vitro* evaluation of buccal tablets of Piroxicam. Int J Pharm Tech Res, 2010; 2(3): 1958-68.
12. Ghosh S, Barik BB. Preparation and evaluation of Aceclofenac sustained release formulation and comparison of formulated and marketed product. Int J Medicine Medical Sci, 2008; 1(9): 375-82.

13. Patel B, Patel P, Bhosalae A, Hardikar S, Mutha S, Chaulang G. Evaluation of tamarind seed polysaccharide as a mucoadhesive and sustained release component of Nifedipine buccoadhesive tablet and comparison with HPMC and Na CMC. Int J Pharm Tech Res, 2009; 1(3): 404-10.
14. Jha KA, Bhattacharya A, Verma PP. Formulation and in vitro evaluation of sustained release matrix tablets of Metoprolol succinate using hydrophilic polymers. Int J Pharm Tech Res, 2009; 1(4): 972-77.
15. Abdelbary GA, Tadros MI. Design and *in vitro/in vivo* evaluation of novel Nicorandil extended release matrix tablets based on hydrophilic interpolymer complexes and a hydrophobic waxy polymer. Eur J Pharm Biopharm, 2008; 69(3): 1019-28.
16. Kulkarni RV, Shah A, Boppana R. Development and evaluation of xyloglucan matrix tablets containing Naproxen. Asian J Pharmaceut, 2008; 2(2): 102-5.
17. Jain S, Yadav SK, Patil UK. Preparation and evaluation of sustained release matrix tablet of Furosemide using natural polymers. Research J Pharm Tech, 2008; 1(4): 374-6.
18. Ganesan V, Jayachandran DL. Design and evaluation of matrix tablets of Ambroxol hydrochloride using guar gum research. J Pharm Tech, 2008; 1(4): 507-12.
19. Punna RR, Sindura G, Ranendra NS. Design and in vitro evaluation of Zidovudine oral controlled release tablets prepared using Hydroxy propyl methy cellulose. Chem. Pharm. Bull, 2008; 56(4): 518-24.
20. Kyeo-re lee, Eun-jung kim, Sang-wan seo, Hoo-kyun choi. Effect of poloxamer on the dissolution of Felodipine and preparation of controlled release matrix tablets containing Felodipine. Arch Pharm Res, 2008; 31(8): 1023-28.
21. Nagaswamy venkatesh D, Sangeetha S, Samantha MK, Shankar SB. Suresh. Design and Evaluation of ethyl cellulose sustained release matrix tablets of Theophylline. International journal of pharmaceutical sciences and Nanotechnology, 2008; 1(1): 60-63.
22. Derle DV, Kasliwal NH , Charan N. Development and comparative evaluation of xanthan gum and Guar Gum based sustained release matrix tablets of Tizanidine hydrochloride. Indian drugs, 2008; 45(6): 485-9.

23. Amelia A, Vikram K. Design and evaluation of matrix based controlled release tablets of Diclofenac Sodium and chondrotin Sulphate. AAPS PharmSciTech, 2007; 8(4): E1-E6.
24. Hamdy A, Ossama YA, Hesham S. Formulation of controlled- release Baclofen matrix tablets: Influence of some hydrophilic polymers on the release rate and In vitro evaluation. AAPS Pharm Sci Tech, 2007; 8(4): E1-E10.
25. Punna RR, Sindhura G, Ranendra NS. Design and study of Lamivudine oral controlled release tablets, AAPS PharmSciTech, 2007; 8(4): E1- E9.
26. Gohel MC, Parikh RK, Padshala MN, Sarvaiya KG, Jena DG. Formulation and optimization of direct compressible isoniazid modified release matrix tablet. Ind J Pharm Sci, 2007; 69.
27. Yeole PG, Galgatte UC, Babla IB, Nakhat PD. Design and evaluation of xanthan gum based sustained release matrix tablets of diclofenac sodium. Ind J Pharm Sci, 2006; 68(2): 185-9.
28. Varshosaz J, Tavakoli N, Kheirolahi F. Use of hydrophilic natural gums in formulation of sustain release matrix tablets of tramadol hydrochloride. AAPS Pharm Sci Tech, 2006; 7(1): 1-8.
29. Senapati MK, Srinatha A, Pandit JK. *In vitro* release characteristics of matrix tablets study of karaya gum and guar gum as release modulators. Ind J Pharm Sci, 2006; 68(6): 824-6.
30. Nerkur J, Jun HW, Park JC, Park MO. Controlled-release matrix tablets of ibuprofen using cellulose ethers and carrageenans: effect of formulation factors on dissolution rates. Eur J Pharm Biopharm, 2005; 61(1-2): 56-68.
31. Saleh M.Al- Saidan, Yellela SRK, Srinivas SP, Vemulapalli S. *In vitro* and *In vivo* evaluation of Guar Gum matrix tablets for oral controlled release of water soluble Diltiazem hydrochloride. AAPS PharmSciTech, 2005; 6(1): E14-E21.
32. Emami J, Tavakoli N, Movahedian A. Formulation of sustained-release lithium carbonate matrix tablets: influence of hydrophilic materials on the release rate and *in vitro-in vivo* evaluation. J Pharm Pharmaceut Sci, 2004; 7(3): 338-44.

33. Hajare AA, More HN, D'Souza JI. Design and evaluation of sustained release tablets of diltiazem hydrochloride. Indian Drugs, 2004; 41(3): 175-6.
34. Shan-yang lin, Mei- jane li , Kung-Hsu lin. Hydrophilic excipients modulate the time lag of time- controlled disintegrating press- coated tablets. AAPS PharmSciTech, 2004; 5(4): 1-5.
35. Crowley MM. Studied Physicochemical properties and mechanism of drug release from ethyl cellulose matrix tablets prepared by direct compression and hot-melt extrusion. International Journal of Pharmaceutics, 2004; 509–522.
36. Koester LS *et al*. Mathematical evaluation of in vitro release profiles of hydroxypropylmethyl cellulose matrix tablets containing carbamazepine associated to b-cyclodextrin. Euro J Pharm and Biopharmaceutics, 2004; 58: 177–179.
37. Reddy KR, Mutalik S, Reddy S. Once-daily sustained-release matrix tablets of nicorandil: formulation and *in vitro* evaluation. AAPS Pharm Sci Tech, 2003; 4(4): 61.
38. Sandip B. Tiwari, T. Krishna murthy, M. Raveendra pai, Pavak R. Mehta , Pasula B. Chowdary. Controlled release formulation of Tramadol hydrochloride using hydrophilic and hydrophobic matrix system. AAPS PharmSciTech, 2003; 4(3): 1-6.
39. Sumathi S, Ray AR. Release behaviour of drugs from tamarind seed polysaccharide tablets. J Pharm pharmaceut Sci, 2002; 5(1): 12-8.
40. Mohan M, Babu GV. Development of new controlled released formulation of flurbiprofen, in-vitro in-vivo correlation. Ind J Pharm Sci, 2002; 64(1): 37-43.
41. Sanchez-Lafuente C, Teresa Faucci M, Fernandez-Arevalo M, lvarez-Fuentes J, Rabasco AM, Mura P. Development of sustained release matrix tablets of didanosine containing methacrylic and ethylcellulose polymers. Int J Pharm, 2002; 234: 213-21.
42. Sanchez-Lafuente C, Furlanetto S, Fernandez-Arevalo M, Alvarez-Fuentes J, Rabasco AM, Faucci MT, Pinzauti S, Mura. Didanosine extended-release matrix tablets: optimization of formulation variables using statistical experimental design. Int J Pharm, 2002; 26: 107-18.
43. Sanchez-Lafuente C, Rabasco AM, Alvarez-Fuentes J, Fernandez-Arevalo M. Eudragit RS-PM and Ethocel 100 Premium: influence over the behavior of didanosine inert matrix system. Farmaco, 2002; 57(8): 649-56.

44. Betageri GV, Deshmukh DV, Gupta RB. Oral sustained-release bioadhesive tablet formulation of didanosine, Drug Dev Ind Pharm. 2001; 27(2): 129-36.
45. Nath BS, Venkatesh, Hiremath D. Formulation and evaluation of sustained release dosage form of theophylline using a combined hydrophobic and hydrophilic matrix, Ind J Pharm Sci. 2000; 62(1): 33-6.
46. Dale L. Munday, Philip JC. Compressed xanthan and Karaya gum matrices: hydration, erosion and drug release mechanisms. International journal of pharmaceutics, 2000; 203: 179-92.
47. Paolo G, Elisabetta G, Mario DLM, Gerolamo P. Evaluation of Alginate compressed matrices as prolonged drug delivery systems. AAPS PharmSciTech, 2000; 1(3): 75-82.
48. Torrado S, Cadorniga R, Torrado JJ. Effects of drug release rate on bioavailablity of different asprin tablets. Int J Pharm, 1996; 133: 65-70.
49. Knupp CA, Milbrath R, Barbhaiya RH. Effect of time of food administration on the bioavailability of didanosine from a chewable tablet formulation. J Clin Pharmacol, 1993; 33(6): 568-73.
50. Basak SC, Jayakumar BM, Mani KPL. Formulation and release behaviour of sustained release ambroxol hydrochloride HPMC matrix tablet. Research paper, Annamalai University: The Madras Pharmaceuticals, Chennai.
51. Nashiru B, Kah HY. Formulation variables affecting drug release from xanthan gum matrices at laboratory scale and pilot scale.
52. Gilbert S, Banker C, Rhodes T (2008), Modern Pharmaceutics. New York, Marcel Dekker Inc, 2008; 4 (121): 503-07.
53. Muralidharan VK, Ganapathy MM, Velayudhan KC, Amalraj VA. Collecting jackfruit germplasm in Western Ghats. Indian J. of Plant Genetic Resources, 1997; 10(2): 227–31.
54. Rahman AM, Nahar N, Mian AJ, Mosihuzzaman M. Variation of carbohydrate composition of two forms of fruit from jack tree (*Artocarpus heterophyllus L.*) with maturity and climatic conditions. Food Chem, 1999; 65: 91-97.
55. Rama Rao AV, Mala Varadan, Venkataraman, Colouring matter of the *A. hetetophyllus*. Indian J. Chem, 1973; 11: 298-299.